

FIFRA Section 2(ee)
Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Crymax[®]
BIOLOGICAL INSECTICIDE

EPA Reg. No. 70051-86

PESTS

For Control of Light Brown Apple Moth (LBAM) (*Epiphyas postvittana*), Banana Moth (*Opogona sacchari*), European Grapevine Moth (*Lobesia botrana*) and European Pepper Moth (*Duponchelia fovealis*).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For Light Brown Apple Moth: Apply when newly hatched larvae appear and before leaves are rolled or webbing is significant.

For Banana Moth: Drench bark to newly emergent shoots following pruning or apply to susceptible plant tissues when Banana Moth larvae are active.

For European Grapevine Moth: Apply at blackhead egg stage or when larvae are newly hatched before leaves are rolled, or larvae have entered fruit.

For European Pepper Moth: Begin applications at egg lay and continue at 3-5 day intervals throughout larval feeding period.

For All Pests: Continue applying as part of a normal spray program until pest is adequately controlled. Apply when caterpillars are actively

feeding. To be effective, Crymax[®] spray must be deposited at the larval feeding site. Crymax can be applied by ground or air in water sufficient to insure thorough and even coverage. Thorough and uniform crop coverage is required for adequate insect control. Applications at higher water volumes have demonstrated improved control of targeted pests. Early morning or evening applications, when air is calm, are generally best for aerial applications.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

Continued on reverse side

Crymax[®]

BIOLOGICAL INSECTICIDE

APPLICATION RATE

Apply Crymax at 0.5 to 2.0 pounds per acre to the following crops:

Vegetable Crops:

Artichokes, Arugala, Asparagus, Beans, Beets, Bok Choy, Broccoli, Brussel Sprouts, Cabbage, Cardoni, Carrots, Cauliflower, Celeriac, Celery, Chick Peas, Chicory, Chinese Cabbage, Collards, Cucumber, Cucurbits, Dry Bulb Onions, Eggplant, Escarole, Endive, Garlic, Green Onions, Greens: (Beets, China, Dandelion, Mustard, Turnip), Horseradish, Kale, Kohlrabi, Leeks, Lentils, Lettuce (Head, Leaf, Romaine), Malanga, Melons (Cantaloupe, Crenshaw, Honeydew, Muskmelon, Watermelon, Etc.), Napa, Okra, Onions, Parsley, Parsnips, Peas, Peppers, Potatoes, Pumpkins, Radishes, Rutabaga, Salsify, Shallots, Soybean Foliage, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips.

Herbs and Spices:

Basil, Chives, Cilantro, Dill, Oregano, Peppermint, Thyme.

Fruit, Nut and Vine Crops:

Pome and Stone Fruits: (Apples, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes, Quince).

Nut Trees: Almonds, Chestnuts, Filberts, Pecans, Pistachios, Walnuts.

Citrus:

Small Fruit and Berries: Blackberries, Blueberries, Boysenberries, Cranberries, Currants, Loganberries, Raspberries, Strawberries.

Grapes:

Tropical and Other Fruit: Avocados, Bananas, Kiwi, Persimmons, Pomegranate, Pineapple, Tropical Fruits.

Field Crops:

Canola/RapeSeed, Evening Primrose, Meadow Foam, Corn (Field, Sweet Popcorn, Seed), Cotton, Hops, Jojoba, Peanuts, Rice, Safflower, Small Grains (Barley, Oats, Rye, Wheat, etc.), Sorghum, Soybeans, Sunflowers, Tobacco.

Commercial Flowers and Ornamental Plants:

Bedding Plants, Flowers (Greenhouse and Field), Greenhouse Ornamentals, Greenhouse Vegetables, Container Stock.

Forest, Shade Tree and Nursery Stock:

Forest Trees, Shade Trees, Nursery Trees