

# Javelin<sup>®</sup>WG

## BIOLOGICAL INSECTICIDE

For Control of Insect Pests of Vegetables, Fruit and Field Crops

 FOR ORGANIC PRODUCTION



**Active Ingredient:**

*Bacillus thuringiensis*, subspecies *kurstaki* strain SA-11 solids, spores and Lepidopteran active toxins\* .....85.0%

**Other Ingredients:** .....15.0%

**TOTAL** .....100.0%

\*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See additional precautionary statements

**FIRST AID**

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Harmful if inhaled. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Personal Protective Equipment:**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow manufacturer's instructions for cleaning/ maintaining PPE.

If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**Net Contents:**

EPA Reg. No. 70051-66  
EPA Est. No. 70051-CA-001  
Lot No.:

Manufactured by  
Certis USA, L.L.C.  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046



**User Safety Recommendations:**

Users should:

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product must not be applied aerially within 1/4 mile of any habitats of endangered or threatened Lepidoptera. No manual application can be made within 300ft. of any threatened or endangered Lepidoptera.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Prosper with pesticides by using them properly! Read and follow label directions.

This labeling must be in the possession of the user at the time of the pesticide application.

### Agricultural Use Requirements:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NOTICE: Read "Limitation of Warranty and Limitation of Liability" on the container or in this Directions for Use section before buying or using. If terms are not acceptable, return at once, unopened.

Unless specifically stated, do not apply this product through any type of irrigation system.

JAVELIN® WG is a biological insecticide for the control of lepidopterous larvae (see Application Rates section).

JAVELIN® WG attacks the larval gut and must be ingested by the insect to be effective.

JAVELIN® WG may be applied up to and on the day of harvest.

### APPLICATION DIRECTIONS

For most consistent control, apply at first sign of newly hatched worms (1st and 2nd instar larvae). Instructions for specific crops are located in the ADDITIONAL INSTRUCTION sections under APPLICATION RATES.

Reapply as necessary under a pest management program that includes close scouting.

If rapid knockdown of heavy worm or non-lepidopterous populations is necessary, include an effective contact insecticide in combination with JAVELIN® WG.

For heavy worm infestations, use the higher JAVELIN® WG rate. During situations of dense foliage and/or rapid growth, increasing water carrier volumes will provide better crop coverage and improve JAVELIN® WG performance.

Tank mix instructions are for use only in states where the tank mix product and application site are registered. Read and follow all label directions for use for other pesticides used as tank mix partners with JAVELIN® WG for specific rates, application timing, and precautions.

### Mixing

Fill spray or mixing tank 3/4 full of water. Turn on agitation and pour JAVELIN® WG into water while maintaining continuous agitation. Add other spray material (if any) and add balance of water. Agitate as necessary to maintain suspension. Do not allow diluted sprays to remain in the tank for more than 48 hours. JAVELIN® WG is formulated to provide desirable coverage and adherence to leaf surfaces. Additional adjuvants, spreaders, or stickers may be added to improve product performance, especially under heavy dew or rainy conditions. Combinations with commonly used insecticides, fungicides, or other spray tank adjuvants are generally not deleterious to JAVELIN® WG if the mix is used promptly. Before mixing in the

spray tank, it is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities.

### Application Instructions

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

### Ground Application

Unless otherwise stated, use the application rate amount of JAVELIN® WG in a minimum of 20 gallons of water per acre depending on type of crop and requirements of state regulations. If lower volumes are used, proper application equipment must be used to insure adequate coverage. Thorough and uniform crop coverage is required for adequate insect control.

### Aerial Application

Use application rate amount of JAVELIN® WG in at least 3 gallons of water per acre. Applications at higher water volumes have demonstrated improved control of targeted pests. Apply early morning or evening when air is calm.

### INSECTS CONTROLLED

When used as directed, JAVELIN® WG will control the following insects.

#### Common Name

Alfalfa caterpillar  
Almond Moth  
Armyworm  
Artichoke plume moth  
Bagworm  
  
Banana moth  
Banana skipper  
Bertha armyworm  
Blueberry leafrollers  
Blueberry spanworm  
Bollworm  
California oak moth  
Cherry fruitworm  
Citrus cutworm  
Codling moth  
Cotton leafperforator  
Cotton leafworm  
Cutworm  
Diamondback moth  
Douglas-fir tussock moth  
  
Elm spanworm  
European corn borer  
European Grapevine moth  
Fall cankerworm  
Fall webworm  
Filbert webworm  
  
Fruittree leafroller  
Grape leafroller  
Grapeleaf skeletonizer  
Green cloverworm  
Green fruitworm  
Gypsy moth  
Helicoverpa spp.  
Heliothis spp.  
Hornworms  
Imported cabbageworm  
Jack pine budworm  
Light brown apple moth  
Loopers

#### Scientific Name

*Colias eurytheme* (Boisduval)  
*Cadra cautella* (Walker)  
*Pseudaletia unipuncta* (Haworth)  
*Platyptilla carduidactyla* (Riley)  
*Thyridopteryx ephemeraeformis* (Haworth)  
*Opogona sacchari*  
*Erionota thrax* (Haworth)  
*Mamestra configurata* various  
*Itame argillacearia* (Pack.)  
*Helicoverpa zea* (Boddie)  
*Phryganidia californica* (Packard)  
*Grapholita packardii* (Zeller)  
*Xylomyges curialis*  
*Cydia pomonella* (Linnaeus)  
*Bucculatrix thurberiella* (Busck)  
*Alabama argillacea* (Hubner)  
various, family Noctuidae  
*Plutella xylostella* (Linnaeus)  
*Orgyia pseudotsugata* (McDunnough)  
*Ennomos subsignaria* (Hubner)  
*Ostrinia nubilalis* (Hubner)  
*Lobesia botrana*\*  
*Alsophila pomataria* (Harris)  
*Hyphantria cunea* (Drury)  
*Melissopus latiferreanus* (Walsingham)  
*Archips argyrospila* (Walker)  
*Desmia funeralis* (Hubner)  
*Harrisina americana* (Guerin)  
*Plathypena scabra* (Fabricius)  
*Lithophane antennata* (Walker)  
*Lymantria dispar* (Linnaeus)  
*Helicoverpa* spp.  
*Heliothis* spp.  
*Manduca* spp.  
*Pieris rapae* (Linnaeus)  
*Choristoneura pinus* (Freeman)  
*Epiphyas postvittana* various

| Common Name              | Scientific Name                              |
|--------------------------|--|
| Mimosa webworm           | <i>Homadaula anisocentra</i> (Meyri)         |
| Naval orangeworm         | <i>Amyelois transitella</i> (Walker)         |
| Obliquebanded leafroller | <i>Choristoneura rosaceana</i> (Harris)      |
| Omnivorous leafroller    | <i>Platynota stultana</i>                    |
| Omnivorous leaf-tier     | <i>Cnephasia longana</i> (Haworth)           |
| Orange tortrix           | <i>Argyrotaenia citrana</i> (Fernald)        |
| Orangedog                | <i>Papilio cressphontes</i> (Cramer)         |
| Oriental fruit moth      | <i>Grapholita molesta</i> (Busck)            |
| Pandemis leafroller      | <i>Pandemis pyrusana</i>                     |
| Peach twig borer         | <i>Anarsia lineatella</i> (Zeller)           |
| Pecan nut casebearer     | <i>Acrobasis nuxvorella</i> (Neunzig)        |
| Redbanded leafroller     | <i>Argyrotaenia velutinana</i> (Walker)      |
| Redhumped caterpillar    | <i>Schizura concinna</i> (J.E. Smith)        |
| Rindworm complex         | various                                      |
| Roughskinned cutworm     | <i>Athetis mindara</i> (Barnes & McDunnough) |
| Saltmarsh caterpillar    | <i>Estigmene acrea</i> (Drury)               |
| Sod webworm              | <i>Crambus mutabilis</i>                     |
| Southwestern corn borer  | <i>Diatraea grandiosella</i> (Dyar)          |
| Spotted cutworm          | <i>Xestia</i> spp.                           |
| Spring cankerworm        | <i>Paleacrita vernata</i> (Peck)             |
| Spruce budworm           | <i>Choristoneura fumiferana</i> (Clemens)    |
| Tent caterpillar         | various, family Lasiocampidae                |
| Tobacco budworm          | <i>Helicoverpa virescens</i> (Fabricius)     |
| Tobacco hornworm         | <i>Manduca sexta</i> (Linnaeus)              |
| Tomato pinworm           | <i>Keiferia lycopersicella</i> (Walsingham)  |
| Tropical sod webworm     | <i>Herpetogramma phaeopteralis</i> (Guenee)  |
| Tufted apple bud moth    | <i>Platynota idaeusalis</i> (Walker)         |
| Variegated leafroller    | <i>Platynota flavedana</i> (Clemens)         |
| Velvetbean caterpillar   | <i>Anticarsia gemmatilis</i> (Hubner)        |
| Western tussock moth     | <i>Orgyia vetusta</i> (Boisduval)            |

\*Apply at blackhead egg stage or when larvae are newly hatched before leaves are rolled or larvae have entered fruit. Continue application as necessary for larval control.

#### RATE SELECTION CONSIDERATIONS

Application rates are typically given as a range:

**Lower rate ranges** may be desired when tank mixing with contact insecticides labeled for worm control or under conditions of light worm infestations or when uniformly small worms are present.

**Medium rate ranges** may be desired when multiple worm life stages are present, continuous egg hatches are occurring or young or light armyworm infestations exist.

**Upper rate ranges** may be desired for heavy worm infestations, mature (larger) worms or for moderate to heavy infestations of armyworm, bollworm, or other difficult to control worm species.

Unless otherwise stated, use the application rate amount of JAVELIN® WG in a minimum of 20 gallons of water per acre depending on type of crop and requirement of state regulations. Lower volumes may be used, but proper application equipment must be used to insure adequate coverage. Thorough and uniform crop coverage is required for adequate insect control.

#### APPLICATION RATES

| Crops  | Lbs./Acre   |
|--|-------------|
| <b>VEGETABLE CROPS, including but not limited to</b>   |             |
| <b>Artichokes</b>  | 0.50 - 1.25 |
| ADDITIONAL INSTRUCTIONS:<br>Apply in a minimum of 100 gal. of water per acre with a spray interval of 10 days or less.<br>Apply 1 lb/A in combination with ASANA® XL, AMBUSH®, or SUPRACIDE® 25-W aerially to aid in resistance management of the artichoke plume moth. Use and follow all label directions of the tank mix partner regarding application, timing, gallonage, and schedules. |             |

| Crops   | Lbs./Acre   |
|---|-------------|
| <b>Asparagus, Beans, (Green, Lima, Mung), Broccoli, Broccoli Raab (Rapini), Brussels Sprouts, Cabbage, Cardoon, Carrots, Cauliflower, Celeriac, Celery, Chick Peas, Chinese Broccoli, Chinese Cabbage, Collards, Cucumbers, Dry Bulb Onions, Eggplants, Garlic, Green Onions, Greens (Dandelion, Turnip, Mustard, Beet, China), Herbs (Basil, Cilantro, Dill, Oregano, Thyme, etc.), Horseradish, Kale, Kohl Rabi, Leeks, Lettuce (Endive, Romaine, Head Lettuce, Escarole, Butter Crunch, Leaf, etc.), Melons (Cantaloupe, Crenshaw, Honeydew, Muskmelon, Watermelon, etc.), Okra, Onions, Parsley, Parsnips, Peas, Peppers, Potatoes, Pumpkins, Radishes, Rutabaga, Salsify, Spinach, Squash (Summer and Winter), Sweet Corn, Sweet Potatoes, Swiss Chard, Table Beets, Tomatoes, Turnip Root, Watercress, Yams</b> | 0.12 - 1.50 |
| ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.   |             |
| <b>FIELD CROPS, including but not limited to</b>  |             |
| <b>Alfalfa (Hay and Seed), Sudan Grass, Hay Crops &amp; Other Forage Crops</b>  | 0.25 - 1.50 |
| ADDITIONAL INSTRUCTIONS:<br>Under conditions of rapid plant growth and rapidly increasing armyworm populations (10 worms or greater per 180° sweep) use the highest rate. Against heterogenous worm populations, where 4th and 5th instars are present, and continuous egg laying is occurring, applications may provide variable control. Under these conditions, the addition of a contact insecticide in combination with JAVELIN® WG is recommended. The addition of a spreader sticker to JAVELIN® WG may provide improved performance.  |             |
| <b>Canola and Evening Primrose</b>  | 0.12 - 1.50 |
| ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.   |             |
| <b>Dry Beans and Peas, Lentils, Mint, Peanuts, Rice, Safflower, Soybeans, Sugar Beets, Sunflower, Sorghum</b>   | 0.25 - 1.50 |
| ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.   |             |
| <b>Field Corn, Pop Corn, Seed Corn</b>  | 0.50 - 1.50 |
| ADDITIONAL INSTRUCTIONS:<br>Make initial application when economically damaging populations exist. Repeat as necessary to maintain control. Applications must be made to early instars prior to entering the ear or plant.  |             |
| <b>Hops</b>   | 0.25 - 1.00 |
| ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.<br>Begin treatment as soon as possible after hatching and before larvae are protected by leaf folds.  |             |

| Crops  | Lbs./Acre   |
|--|-------------|
| <b>Jojoba</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply in a minimum of 50 gallons of water per acre by ground equipment or a minimum of 10 gallons of water by aerial equipment. Thorough coverage of foliage is essential and dictates the minimum spray volumes necessary.   | 0.50 - 1.00 |
| <b>Small Grains</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.   | 1.00 - 1.50 |
| <b>Tobacco</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.  | 0.12 - 1.25 |
| <b>Cotton</b><br><b>Including Arizona and California</b><br>Early and Mid-Season<br>ADDITIONAL INSTRUCTIONS:<br>Repeat as necessary throughout season to maintain control. If egg laying frequency indicates future moderate to heavy worm populations, time application spray to coincide with the 2nd instar larvae. During periods of high temperatures, worms will progress through 1st and 3rd instars very rapidly and early application timing is necessary for control.<br><br>To be effective, JAVELIN® WG spray must be deposited at the larval feeding site. When plant cover is dense and worms are feeding in the lower 2/3 portion of the plant, aerial application of JAVELIN® WG may not provide adequate control.<br><br>For the suppression of light to moderate infestations, apply at first sign of egg-laying or newly-hatched worms (1st instar larvae). | 0.50 - 1.50 |
| <b>Cotton</b><br><b>Except Arizona and California</b><br>Early<br>ADDITIONAL INSTRUCTIONS:<br>For early season management of <i>Helicoverpa</i> and <i>Heliothis</i> species. Initiate applications when 50% of plants are at pinhead square cotton stage, independent of <i>Helicoverpa</i> and <i>Heliothis</i> egg and larval counts, or at 1st egg lay, whichever occurs earlier. Continue applications on 5-day spray interval up to synthetic pyrethroid spray window. For added control of <i>Helicoverpa</i> and <i>Heliothis</i> , tank mixing of JAVELIN® WG with a labeled ovicide, such as, methomyl (0.125 lb. a.i./acre), profenofos (0.25 lb. a.i./acre), or thiodicarb (0.125-0.25 lb.a. i./acre) is recommended.<br><br>Read and follow all directions for use, precautions and restrictions on tank mix product labels.                                      | 0.25 - 1.25 |
| <b>FRUIT, NUT &amp; VINE CROPS, including but not limited to</b>   |             |
| <b>Apples and Pears</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply when newly hatched larvae appear and before leaves are rolled. Continue applying as a part of the normal cover spray program until pest is adequately controlled. Apply when caterpillars are actively feeding (2nd-4th instars).   | 0.50 - 4.00 |

| Crops   | Lbs./Acre   |
|---|-------------|
| <b>Avocados</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control. Begin treatment as soon as possible after hatching and before larvae are protected by leaf folds.<br>(Amorbia [Mexican leafroller] is suppressed only).  | 0.50 - 1.25 |
| <b>Bananas</b><br>ADDITIONAL INSTRUCTIONS:<br>Hawaii only. Use calibrated ground equipment with adequate water to apply to point of runoff.   | 0.50 - 1.00 |
| <b>Citrus</b><br>ADDITIONAL INSTRUCTIONS:<br>Use 50-600 gallons of water per acre when using ground equipment and 10 gallons of water minimum per acre by air.(Amorbia [Mexican leafroller] is suppressed only.)  | 0.25 - 1.50 |
| <b>Coffee</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.  | 0.25 - 4.00 |
| <b>Blueberries, Caneberries, Currants, Kiwi</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply by ground or aerial equipment using enough water to provide adequate coverage. Begin treatment as soon as possible after hatching. For leafrollers, apply before larvae are protected by leaf folds.   | 0.25 - 1.00 |
| <b>Grapes</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply by ground equipment in up to 200 gallons total spray per acre to obtain thorough coverage of leaf surfaces. Start treating as soon as possible after hatching and before larvae are protected by leaf folds.   | 0.50 - 1.25 |
| <b>Almonds, Apricots, Cherries, Filberts, Nectarines, Peaches, Pecans, Persimmons, Plums, Pomegranates, Prunes, Walnuts</b><br>ADDITIONAL INSTRUCTIONS:<br>For leafrollers, start treating as soon as possible after hatching and before larvae are protected by leaf folds.<br><br>Apply when caterpillars are actively feeding (2nd-4th instar).<br><br>Application timing is very important for good casebearer suppression. Consult your local university or extension agent for information concerning specific modeling, that predicts egg lay, typical application dates, and scouting techniques for your area. JAVELIN® WG must be present at egg hatch for best control. Make application when the majority of eggs are in the pink stage. For best control make two applications 7 days apart. If only one application is made, a minimum of 1lb. should be applied. | 0.25 - 4.00 |

| Crops  | Lbs./Acre   |
|--|-------------|
| <b>Melons (Also see vegetables)</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply at first sign of hatch before larvae enter fruit. Repeat as necessary to maintain control.  | 0.50 - 0.75 |
| <b>Strawberries</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control. Use 20 gallons water minimum per acre when using ground equipment and 5 gallons water minimum per acre by aircraft.<br><br>In a tank mix with contact insecticides, rates as low as 1/2 lb. of JAVELIN® WG may be used for the control of armyworm. | 0.25 - 1.50 |

|  |             |
|--|-------------|
| <b>Olives</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control. | 0.25 - 4.00 |
|--|-------------|

|   |             |
|---|-------------|
| <b>Tropical/Sub-Tropical Fruit</b><br><b>Lychee, Sugar Apple, Papaya, Guava</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply as necessary to maintain control.<br>Begin treatment as soon as possible after hatching and before larvae are protected by leaf folds. | 0.25 - 4.00 |
|---|-------------|

|  |             |
|--|-------------|
| <b>SHADE TREES AND ORNAMENTALS</b><br><b>(Including Roses)</b><br>ADDITIONAL INSTRUCTIONS:<br>Apply when leaf expansion reaches 40% to 50% as infestation warrants.<br>If eggs hatch over a long period of time, or if reinfestation occurs, spray about 14 days after first application.<br><br>Apply when most larvae are 3rd-4th instar. Also consider the opening of the bud cap to ensure foliage exposure. Apply after eggs have hatched and early instar larvae are feeding on exposed foliage. | 0.12 - 1.25 |
|--|-------------|

|   |      |
|---|------|
| <b>TURF AND GRASS SEED PRODUCTION</b><br>ADDITIONAL INSTRUCTIONS:<br>Repeat as necessary throughout season to maintain control. | 1.00 |
|---|------|

**STORED SOYBEANS, GRAINS**  
**(Indian Meal Moth, Almond Moth)**

ADDITIONAL INSTRUCTIONS:  
To control and prevent Indian Meal Moth and Almond Moth infestations of stored soybeans and grains, prepare a spray mixture which includes 1 gallon of water for every 1.5 oz. by weight of JAVELIN® WG. The spray mixture may be applied either by treating the top 4 inches of grain as it is being augered into storage (applying 0.6 pint of mixture per bushel in the grain stream), or by treating the surface of grain after it is in the bin. The Table below can be used as a guide in determining the total amount of JAVELIN® WG needed according to bin diameter or the number of bushels to be treated.

| Bin Diameter (ft.) | Surface Area (sq. ft.) | Bushels (to 4 in. depth) | JAVELIN® WG Rate (by weight) |       |
|--------------------|------------------------|--------------------------|------------------------------|-------|
|                    |                        |                          | Grams                        | oz.   |
| 8                  | 50                     | 13                       | 21                           | 0.75  |
| 12                 | 113                    | 30                       | 50                           | 1.75  |
| 16                 | 201                    | 53                       | 85                           | 3.00  |
| 20                 | 314                    | 84                       | 120                          | 4.25  |
| 24                 | 452                    | 120                      | 185                          | 6.50  |
| 28                 | 615                    | 163                      | 255                          | 9.00  |
| 32                 | 804                    | 214                      | 326                          | 11.50 |

To insure thorough coverage when making applications to the grain surface after it is in the bin, apply spray mixture in three (3) applications. Mix the grain with a scoop or rake to a depth of four (4) inches after each application.

Stored grain may be treated anytime, but for best results, treat grain at the time it is placed into storage or shortly thereafter, or in the early spring prior to egg-laying. Full season control is normally experienced. Re-treat only if reinfestation occurs.

For the protection of bagged grain, apply spray mixture to entire grain mass, and mix thoroughly prior to bagging. JAVELIN® WG at 6 oz. by weight per 10 gallons of water will treat approximately 100 bushels.

Treated grain may be used at any time after treatment.

**Flowers and Ornamentals**

JAVELIN® WG may also be used on flowers and ornamentals outdoors and in the greenhouse at a rate of 0.25-1.50 lb. per 100 gallons of water for control of listed insects on this label.

**Guide for Small Spray Volume Mixing**

| Rate<br>Lbs. / A | Conversion Rate*<br>Teaspoons / Gallon |
|------------------|--|
| 1/4              | 1/2                                    |
| 1/2              | 1                                      |
| 1                | 2                                      |

\*Assumes Application to spray runoff.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**Storage:** Store in original container in a cool, dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing. Storage at temperatures above 90° F may impair effectiveness.

**Pesticide Disposal:** Pesticide, spray mixture, or rinse water that cannot be used according to label instruction must be disposed of according to Federal, State, or Local procedures.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY**

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

**REGISTERED TRADEMARKS**

Ambush® is a registered trademark of AMVAC

Asana® XL is a registered trademark of E.I duPont de Nemours and Co., Inc.

Supracide® 25-W is a registered trademark of Gowan Company.

JAVELIN® WG is a registered trademark of Certis USA, L.L.C.