

SUPPLEMENTAL LABELING:  
HEMP

# LifeGard® WG

BIOLOGICAL PLANT ACTIVATOR



CAN BE USED IN ORGANIC PRODUCTION



**ACTIVE INGREDIENT:**

*Bacillus mycoides* isolate J\* ..... 40.0%

**OTHER INGREDIENTS:** ..... 60.0%

**TOTAL** ..... 100.0%

\* Equivalent to a minimum of 30 billion ( $3 \times 10^{10}$ ) viable spores/g of product.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

This Supplemental Labeling expires on November 1, 2021 and must not be used or distributed after this date.

**MANUFACTURED BY:**

Certis USA LLC  
9145 Guilford Road, Suite 175  
Columbia, MD 21046

**EPA Reg. No. 70051-119  
EPA Est. No. 70051-CA-001**



This is a Specimen Label. It may not reflect the most-recent approved label for use in your state. Always refer to the label on the product packaging for approved use instructions. Please contact your Certis sales representative for more information.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for LifeGard® before applying.

Use of LifeGard® according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for LifeGard®.

## **CROP**

Hemp

## **PEST**

Anthrachnose (*Colletotrichum* spp.), Brown blight (*Alternaria alternata*), Brown leaf spot and stem canker (*Ascochyta* spp.), Gray mold (*Botrytis cinerea*), Hemp leaf spot (*Bipolaris* sp.), Powdery mildew (*Leveillula*, *Podosphaera*, *Sphaerotheca* spp.), White leaf spot (*Phomopsis ganjiae*), Yellow leaf spot (*Septoria* spp.), Olive leaf spot (*Cercospora cannabina*), Stemphylium leaf and stem spot (*Stemphylium botryosum*), Bacterial blight (*Pseudomonas cannabina*), Xanthomonas leaf spot (*Xanthomonas campestris*)

## **RATE**

Apply LifeGard® at a concentration of 4.5 ounces (128 grams) per 100 gallons of water. If using dry measure rather than weight, the volume of 4.5 ounces of LifeGard® is approximately  $\frac{3}{4}$  cup.

## **APPLICATION METHOD**

Apply at first appearance of leaves or just after transplant and repeat at 3-14 day intervals as needed, in sufficient water to obtain thorough coverage of foliage.

*Ground applications:* LifeGard® can be applied in most commonly-used ground application equipment, such as tractor-mounted boom, airblast, high clearance, backpack, and other pressurized sprayers; hose-end or hand-held sprayers; and foggers or mist blowers. Apply in sufficient volume of water to provide uniform coverage.

*Aerial applications:* LifeGard® can be applied by fixed or rotary winged aircraft in a minimum of 5 gallons of water per acre. Standard precautions should be taken to minimize spray drift.

*Chemigation:* LifeGard® can be applied to the crop canopy through overhead sprinkler systems by injecting the specified rate at the very end of the irrigation period. Injection should occur only within the minimum time required to ensure complete flushing of the product from the system and onto the crop canopy. Keep supply tank agitated during application. See "Chemigation Instructions" on label affixed to the container for LifeGard® for additional information. Do not apply LifeGard® through any other type of irrigation system.

ESL20200427

Ver20200428