

**FIFRA Section 2(ee) Recommendation  
For Control of Tomato Bacterial Canker on  
Fruiting Vegetables (Crop Group 8-10)  
FOR USE AND DISTRIBUTION IN THE STATE OF NY ONLY**

***This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.***

**All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed.** Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

***Listed Crops***

Pathogen	Application Rate	Use Instructions
Tomato Bacterial Canker ( <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> )	Foliar: 1 to 3 quarts per acre	Tank mix or rotate with copper-based fungicides at label rates for improved control.

**Crop-Specific Restrictions:**

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

TO ENSURE UNIFORMITY, MIX PRODUCT WELL BEFORE USE.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

